



2016 State of

# Drunk Driving Fatalities IN AMERICA

#### **BOARD OF DIRECTORS**

Bacardi U.S.A., Inc.

Beam Suntory Inc.

**Brown-Forman** 

Constellation Brands, Inc.

**DIAGEO** 

**Edrington** 

Mast-Jägermeister US, Inc.

Hood River Distillers, Inc.

**Pernod Ricard USA** 

#### **NATIONAL ADVISORY BOARD**

#### The Honorable Thurbert E. Baker

Partner, Dentons US Former Georgia Attorney General

#### **Steven Casstevens**

Chief of Police, Buffalo Grove, Illinois 2nd Vice President of the IACP

#### Charles G. Curie

Principal, The Curie Group, LLC Former Administrator, Substance Abuse and Mental Health Services Administration

#### Jeannette Kaplun

Chief Executive Officer, Hispana Global TV host and author

#### Deborah Gilboa, M.D.

Clinical Associate Professor at University of Pittsburgh, School of Medicine Parenting and Youth Development Expert

#### Lisa Graham Keegan

Principal, Keegan Company Former Superintendent of Public Instruction State of Arizona

#### **Robin Roberts**

Co-founder and President, National Media Group


#### Dean Wilkerson, JD, MBA, CAE

Executive Director, American College of Emergency Physicians

## Fatal Motor Vehicle Crashes in 2016 Included a Variety of Factors




Drunk, drugged, distracted, drowsy, or a combination are forms of impaired driving and are dangerous and preventable.



### **Saving Lives for Over 25 Years**



Since the inception of Responsibility.org in 1991, nearly 90,000 lives have been saved.




Source: NHTSA, FARS, 10/17

### Motor Vehicle Fatalities: 1982-2016



Drunk driving fatalities are descending at a faster rate than total motor vehicle fatalities.




Source: NHTSA, FARS, 10/17

### Percent of 2016 Driving Fatalities by BAC Test Result



In 2016, drunk driving accounted for 28% of all traffic fatalities. Nearly sixty percent of alcoholinvolved fatal crashes involved high BAC drivers.






Source: NHTSA, FARS ARF of Drivers with Known BAC, 10/17 Total may not equal the sum of categories due to rounding.

## Alcohol-Impaired Driving Fatality Data 2016






Source: NHTSA, FARS, 10/17 Total may not equal sum of the states due to rounding. **US TOTAL 10,497** 

## Under 21 Alcohol-Impaired Driving Fatality Data 2016






Source: NHTSA, FARS, 10/17 Total may not equal sum of the states due to rounding. **US TOTAL 1,031** 

### Percent Change in Alcohol-Impaired Driving Fatalities from 2015 to 2016






Source: NHTSA, FARS, 10/17

# Under 21 Percent Change in Alcohol-Impaired Driving Fatalities from 2015 to 2016





Source: NHTSA, FARS, 10/17


### Alcohol-Impaired Driving Fatalities 1982-2016



#### TOTAL ALCOHOL-IMPAIRED DRIVING FATALITIES



### RATE OF TOTAL ALCOHOL-IMPAIRED DRIVING FATALITIES PER 100,000 POPULATION



Top Chart - Source: NHTSA/FARS, 10/17 Bottom Chart - Source: NHTSA, FARS and U.S. Census Bureau, 10/17


### Under 21 Alcohol-Impaired Driving Fatalities 1982-2016



#### **UNDER 21 ALCOHOL-IMPAIRED DRIVING FATALITIES**




### RATE OF UNDER 21 ALCOHOL-IMPAIRED DRIVING FATALITIES PER 100,000 POPULATION



Top Chart - Source: NHTSA/FARS, 10/17 Bottom Chart - Source: NHTSA, FARS and U.S. Census Bureau, 10/17

# 2016 Alcohol-Impaired Driving Fatalities Per 100,000 Population by State





In 22 states and D.C., alcohol-impaired driving fatalities per 100,000 population were at or below the national average of 3.3 deaths per 100,000 population.


Alabama	5.8
Alaska	4.2
Arizona	3.4
Arkansas	3.9
California	2.7
Colorado	2.9
Connecticut	2.8
Delaware	3.9
D.C.	1.5
Florida	4.1
Georgia	3.6
Hawaii	2.5
Idaho	4.6
Illinois	2.5
Indiana	3.2
lowa	3.4
Kansas	3.3

Kentucky	4.0
Louisiana	4.8
Maine	4.1
Maryland	2.2
Massachusetts	1.7
Michigan	2.4
Minnesota	1.7
Mississippi	4.3
Missouri	4.0
Montana	8.2
Nebraska	3.3
Nevada	3.4
New Hampshire	3.0
New Jersey	1.5
New Mexico	5.7
New York	1.4
North Carolina	3.5
North Dakota	6.7

)hio	2.8
)klahoma	4.6
)regon	3.8
Pennsylvania	2.6
Rhode Island	1.8
outh Carolina	6.7
outh Dakota	5.3
ennessee	3.4
exas	5.2
Jtah	1.7
ermont (	4.3
'irginia	2.7
Vashington	2.2
Vest Virginia	3.7
Visconsin	3.3
Vyoming	5.5
JS Total	3.3

# 2016 Under 21 Alcohol-Impaired Driving Fatalities Per 100,000 Population by State





In 26 states and D.C., under 21 alcohol-impaired driving fatalities per 100,000 population were at or below the national average of 1.2 deaths per 100,000 population.


Alabama	2.7
Alaska	1.4
Arizona	1.6
Arkansas	1.2
California	1.3
Colorado	1.2
Connecticut	0.9
Delaware	1.7
D.C.	0.0
Florida	1.6
Georgia	1.0
Hawaii	1.4
Idaho	1.4
Illinois	0.7
Indiana	1.1
lowa	0.8
Kansas	0.8

Kentucky	0.9
Louisiana	1.3
Maine	2.3
Maryland	0.6
Massachusetts	1.0
Michigan	1.2
Minnesota	0.3
Mississippi	1.8
Missouri	1.7
Montana	3.0
Nebraska	1.3
Nevada	1.2
New Hampshire	0.6
New Jersey	0.3
New Mexico	2.6
New York	0.6
North Carolina	1.5
North Dakota	2.4

Ohio	0.9
Oklahoma	1.3
Oregon	1.5
Pennsylvania	0.8
Rhode Island	0.8
South Carolina	1.8
South Dakota	2.8
Tennessee	0.6
Texas	1.9
Utah	0.5
Vermont	0.7
Virginia	1.1
Washington	0.7
West Virginia	1.1
Wisconsin	1.3
Wyoming	1.9
US Total	1.2


# Percent Change in Drunk Driving Fatality Rates from 2007 to 2016 by State Rank






# Percent Change in Under 21 Drunk Driving Fatality Rates from 2007 to 2016 by State Rank





## 2016 Fatal Drunk Driving Crashes Among Drivers with a Known BAC





drivers involved in fatal drunk driving crashes had a BAC level of


Percent of Alcohol-Impaired Driving Fatalities by BAC

Source: NHTSA, FARS ARF of Drivers with Known BAC, 10/17

## Repeat Offenders Involvement in Fatal Crashes



One third of drivers involved in a fatal crash who had a prior DWI\* in the past five years had a high BAC at the time of the crash.




<sup>\*</sup> Prior convictions only for events occurring within 5 years from date of crash. Total may not equal sum of categories due to rounding.

### **Holiday Traffic Fatalities by BAC Level**



Drunk driving fatalities accounted for 36% of traffic fatalities during the 2016 holidays and drivers with a high BAC were involved in 68% of drunk driving fatalities.



The number of whole days in the holiday period depends on the day on which the legal holiday falls. Typically, the period is 6 p.m. the night before until 5:59 a.m. the day after the holiday. Total may not equal sum of categories due to rounding.

### **Terminology**



#### **Alcohol-Impaired Fatality**

Drivers in all 50 states and D.C. are considered to be alcohol-impaired if their blood alcohol concentration (BAC) is .08 grams per deciliter (g/dL) or higher. Any fatality occurring in a crash involving at least one driver, or motorcycle operator, with a BAC of .08 or higher is considered to be an alcohol-impaired driving, or drunk driving, fatality. The term alcohol-impaired does not indicate that a crash or a fatality was the result of alcohol impairment. In producing national and state alcoholimpaired statistics, the National Highway Traffic Safety Administration (NHTSA) estimates the extent of alcohol involvement when alcohol test results are unknown.

#### Alcohol-Involved Fatality

Alcohol-involved fatalities are those where at least one driver, or motorcycle operator, has a positive BAC of .01 or higher. In producing national and state alcoholinvolved statistics, NHTSA estimates the extent of alcohol involvement when alcohol test results are unknown.

#### **Blood Alcohol Concentration (BAC)**

BAC is measured in grams of alcohol per 100 milliliters of blood. A BAC of .01 indicates .01 grams of alcohol per 100 milliliters of blood. As of July 2004, all 50 states and D.C. have passed legislation establishing a driver with a BAC of .08 to be legally intoxicated. Additionally, 48 states and D.C. have laws that increase penalties for those convicted of driving with elevated or "high" BAC levels.

#### Rates per 100,000 Population

The rate of alcohol-impaired fatalities per 100,000 population is the number of alcohol-impaired fatalities for every 100,000 persons in the population being measured. For example, an alcohol-impaired fatality rate of 3.3 per 100,000 population nationally means that for every 100,000 people in the nation, there were approximately three alcohol-impaired fatalities.

#### Repeat Offenders

The NHTSA/FARS data records prior driving records (convictions only, not violations) for driving while intoxicated events occurring within five years of the date of the crash. The same driver can have one or more of these convictions during this five year period. Drivers who have a prior conviction in this five year period are reported as repeat offenders.

The information presented in this report is drawn from several databases maintained by government agencies, including the National Center for Statistics and Analysis (NCSA) of NHTSA, which compiles crash data from the Fatality Analysis Reporting System (FARS), and the U.S. Census Bureau of the U.S. Department of Commerce. This report reflects data from 1982 to the present utilizing NHTSA's multiple imputation method for estimating missing information about BAC levels for persons involved in fatal crashes, thus allowing for improved reporting of alcohol involvement statistics at any BAC level. The U.S. Census Bureau publishes state resident population estimates which were used in this report.





responsibility.org/blog



twitter.com/goFAAR



facebook.com /goFAAR



youtube.com/goFAAR



instagram.com/go\_FAAR



flickr.com/goFAAR



pinterest.com/goFAAR



responsibility.org/get-the-facts



















2345 Crystal Drive Suite 710 Arlington, VA 22202 202-637-0077

RESPONSIBILITY.ORG